

Crystallization and preliminary X-ray diffraction analysis of the lactonase *VmoLac* from *Vulcanisaeta moutnovskia*


Julien Hiblot, Guillaume Gotthard, Charlotte Champion, Eric Chabriere and Mikael Elias

Acta Cryst. (2013). **F69**, 1235–1238

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Republication of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>



Acta Crystallographica Section F: Structural Biology and Crystallization Communications is a rapid all-electronic journal, which provides a home for short communications on the crystallization and structure of biological macromolecules. Structures determined through structural genomics initiatives or from iterative studies such as those used in the pharmaceutical industry are particularly welcomed. Articles are available online when ready, making publication as fast as possible, and include unlimited free colour illustrations, movies and other enhancements. The editorial process is completely electronic with respect to deposition, submission, refereeing and publication.

Crystallography Journals **Online** is available from journals.iucr.org

Julien Hiblot,^{a‡} Guillaume
Gotthard,^{a‡} Charlotte
Champion,^a Eric Chabriere^{a*} and
Mikael Elias^{b*}

^aUnité de Recherche sur les Maladies
Infectieuses et Tropicales Emergentes (URMITE),
UMR CNRS–IRD–INSERM, IHU Méditerranée
Infection, Aix–Marseille Université, 7 Boulevard
Jean Moulin, 13385 Marseille CEDEX 05,
France, and ^bBiological Chemistry, Weizmann
Institute of Science, Rehovot, Israel

‡ These authors made equal contributions.

Correspondence e-mail:
eric.chabriere@univmed.fr,
mikael.elias@weizmann.ac.il

Received 29 July 2013
Accepted 5 September 2013



© 2013 International Union of Crystallography
All rights reserved

Crystallization and preliminary X-ray diffraction analysis of the lactonase *VmoLac* from *Vulcanisaeta moutnovskia*

Phosphotriesterase-like lactonases (PLLs) are native lactonases that are capable of hydrolyzing lactones such as aliphatic lactones or acyl-homoserine lactones, which are involved in bacterial quorum sensing. Previously characterized PLLs are moreover endowed with a promiscuous phosphotriesterase activity and are therefore able to detoxify organophosphate insecticides. A novel PLL representative, dubbed *VmoLac*, has been identified from the hyperthermophilic crenarchaeon *Vulcanisaeta moutnovskia*. Because of its intrinsic high thermal stability, *VmoLac* may constitute an appealing candidate for engineering studies with the aim of producing an efficient biodecontaminant for organophosphorus compounds and a bacterial antivirulence agent. In combination with biochemical studies, structural information will allow the identification of the residues involved in substrate specificity and an understanding of the enzymatic catalytic mechanisms. Here, the expression, purification, crystallization and X-ray data collection at 2.4 Å resolution of *VmoLac* are reported.

1. Introduction

Organophosphates (OPs) are well known neurotoxic compounds that irreversibly inhibit acetylcholinesterase, a key enzyme of the central nervous system (Masson *et al.*, 2009). These compounds, which are massively used as pesticides, are responsible for soil and water pollution, for which no satisfactory means of remediation are available (Singh, 2009). Indeed, existing methods for removing them are cost-prohibitive and cause environmental concerns (LeJeune *et al.*, 1998). Moreover, before World War 2 these compounds were engineered as chemical warfare agents and were used in the Iran–Iraq war and in the terrorist attack on the Tokyo subway (Gupta, 2009). The use of enzymes that are capable of hydrolyzing these compounds represents an appealing alternative to chemical methods (Singh, 2009). Several organophosphate hydrolases (OPHs) belonging to different enzyme superfamilies have been identified and studied [*e.g.* organophosphate acid anhydrolase (OPAA; Vyas *et al.*, 2010), OPHC2 (Gotthard *et al.*, 2013) and paraoxonases (PONs; Ben-David *et al.*, 2012)]. The most studied OPHs comprise the bacterial phosphotriesterases (PTEs) isolated from *Brevundimonas diminuta* (*BdPTE*; Benning *et al.*, 1994) and *Agrobacterium radiobacter* (*OpdA*; Jackson *et al.*, 2006). These enzymes, which exhibit near-diffusion-limit kinetic rates against the insecticide paraoxon (*i.e.* $k_{\text{cat}}/K_{\text{M}} \simeq 10^8 \text{ M}^{-1} \text{ s}^{-1}$; Omburo *et al.*, 1992), are believed to have emerged from the first uses of OPs as insecticides in the 1950s.

Recently, a protein family sharing ~30% sequence identity with PTEs was identified by virtue of their ability to hydrolyze insecticides and were initially dubbed paraoxonases (Merone *et al.*, 2005). A detailed biochemical and phylogenetic analysis later revealed that these proteins, named phosphotriesterase-like lactonases (PLLs), are in fact native lactonases endowed with promiscuous phosphotriesterase activity (Afriat *et al.*, 2006; Elias & Tawfik, 2012). PLLs are likely to be the progenitors of PTEs (Afriat-Jurnou *et al.*, 2012). PLLs hydrolyze various lactones, including aliphatic lactones and acyl-homoserine lactones (AHLs), that are involved in bacterial quorum sensing (Hiblot *et al.*, 2012a; Hawwa, Larsen *et al.*, 2009). The ability of lactonases such as PLLs to hydrolyze AHLs enables these enzymes

to interfere with bacterial communication (Dong *et al.*, 2001) and they therefore offer interesting potentialities to develop new approaches in order to fight against several pathogens (Amara *et al.*, 2011).

PTEs and PLLs, which belong to the amidohydrolase superfamily (Seibert & Raushel, 2005), share the same $(\alpha/\beta)_8$ topology in which a bimetallic centre is coordinated by four histidines, an aspartic acid and a carboxylated lysine (Elias *et al.*, 2008; Del Vecchio *et al.*, 2009). The bimetallic centre activates a water molecule into a hydroxide ion, which serves as a nucleophile for the hydrolysis of OPs or AHLs. The catalytic centre is surrounded by two loops involved in the substrate specificity: loops 7 and 8 (Elias *et al.*, 2008; Jackson *et al.*, 2009). PTEs and PLLs differ mainly in the relative size and conformations of these loops, which account for the different substrate specificity of these enzymes (Afriat-Jurnou *et al.*, 2012). PLLs hydrolyze organophosphate compounds with poor to moderate efficiency (Hiblot *et al.*, 2012*b*; Hawwa, Larsen *et al.*, 2009; Zhang *et al.*, 2012) compared with PTEs (Omburo *et al.*, 1992; Jackson *et al.*, 2006; Donarski *et al.*, 1989). Some PLL representatives, however, offer interesting biotechnological potentialities for engineering an efficient organophosphate biodecontaminant because of their high thermal stability (Hiblot *et al.*, 2012*a,b*; Hawwa, Larsen *et al.*, 2009; Hawwa, Aikens *et al.*, 2009), as illustrated by several engineering studies on these enzymes (Merone *et al.*, 2010; Hawwa, Larsen *et al.*, 2009; Xue *et al.*, 2013; Chow *et al.*, 2010).

VmoLac (YP_004245953) is an enzyme identified from the recently sequenced crenarchaeon *Vulcanisaeta moutnovskia* strain 768-28 (Gumerov *et al.*, 2011). This organism was isolated from solfataric fields close to the Moutnovsky volcano in Kamchatka, Russia. Its growth temperature ranges between 333 and 371 K (Gumerov *et al.*, 2011). *VmoLac* shares ~50% sequence identity with other PLLs. Here, we report the protein production, purification, crystallization and preliminary X-ray diffraction of *VmoLac*.

2. Cloning, expression and purification of *VmoLac*

The gene encoding *VmoLac* (YP_004245953) was optimized for *Escherichia coli* expression by the GeneArt service provider (Life Technologies, France). The optimized gene included N-terminal tags [a *Strep*-tag (MSAWSHPQFEK) for affinity chromatography purification and a TEV cleavage site (ENLYFQ/G) for removal of the tag; Gotthard *et al.*, 2011] and was synthesized by GeneArt (Life Technologies, France). This construct leaves an N-terminal Gly residue after cleavage of the tag by TEV protease. The complete gene was subsequently cloned by the same provider into a custom version of pET-22b(+) (Novagen) using *NdeI* and *XhoI* as cloning sites. Recombinant *VmoLac* protein was overexpressed using a protocol that was previously used for another PLL (Gotthard *et al.*, 2013; Hiblot *et al.*, 2012*a*). Briefly, recombinant *VmoLac* protein was overproduced in *E. coli* BL21 (DE3)-pGro7/GroEL strain (TaKaRa). Protein expression was performed in 4 l ZYP medium (100 $\mu\text{g ml}^{-1}$ ampicillin, 34 $\mu\text{g ml}^{-1}$ chloramphenicol) inoculated with a 50 ml overnight pre-culture. The culture was grown at 310 K until the $\text{OD}_{600\text{ nm}}$ reached 0.6. Induction of the protein was conducted by consumption of the lactose in the ZYP medium, a temperature transition to 298 K over 20 h and the addition of 0.2 mM CoCl_2 .

The cells were harvested by centrifugation (4500g, 277 K, 15 min). The pellets were resuspended in lysis buffer [50 mM HEPES pH 8, 150 mM NaCl, 0.2 mM CoCl_2 , 0.25 mg ml^{-1} lysozyme, 10 $\mu\text{g ml}^{-1}$ DNase I, 0.1 mM phenylmethylsulfonyl fluoride (PMSF)] and stored at 193 K for 2 h. The suspended frozen cells were thawed at 310 K for 15 min and disrupted by three sonication steps of 30 s (Branson

Sonifier 450; 80% intensity and micro tip limit at 8). Cell debris was removed by centrifugation (14 500g, 277 K, 30 min). The cell lysate was then loaded onto a *StrepTrap* column (GE Healthcare) at a flow rate of 5 ml min^{-1} and protein elution was performed using elution buffer (50 mM HEPES pH 8, 150 mM NaCl, 0.2 mM CoCl_2 , 2.5 mM desthiobiotin). Because of the low binding capacity of the column, this step was repeated five times and all of the protein was then cleaved by TEV protease (1250 μg , 20 h, 310 K; van den Berg *et al.*, 2006). Spontaneously precipitated TEV protease was harvested by centrifugation (12 000g, 277 K, 10 min). The protein was subsequently concentrated using a centrifugation device (Amicon Ultra MWCO 10 kDa; Millipore, Ireland) prior to a size-exclusion chromatography step (Superdex 75 16/60, GE Healthcare) in 50 mM HEPES pH 8, 150 mM NaCl, 0.2 mM CoCl_2 . Fractions containing pure protein were pooled and concentrated prior to crystallization trials using a centrifugation device (Amicon Ultra MWCO 10 kDa; Millipore, Ireland). The yield of production was about 5 mg per litre of culture.

The purity of the protein was checked with Coomassie-stained 15% SDS-PAGE (Fig. 1), which revealed two main bands (35 and 70 kDa). Both bands were subjected to mass-spectrometric analysis (MS Platform Timone, Marseille, France) and both were identified as the *VmoLac* protein. The molecular mass of the *VmoLac* monomer being 35 548 Da, the two bands on the gel at 35 and 70 kDa were attributed to monomers and dimers of *VmoLac*, respectively. As the *VmoLac* dimer originates from an extremely thermophilic organism, its dimer may resist the denaturing conditions of sample preparation for SDS-PAGE experiments (368 K incubation for 10 min, 715 mM β -mercaptoethanol).

3. Protein crystallization

VmoLac was concentrated to 20 mg ml^{-1} for crystallization trials. Crystallization assays were performed using the sitting-drop vapour-diffusion method setup in a 96-well plate and the commercial screen conditions Structure Screen 1 + 2 (Molecular Dimensions). The plate was incubated at 277 K. Crystals appeared after 1 d at 277 K in a




Figure 1
15% SDS-PAGE of *VmoLac* protein performed under denaturing conditions and stained with Coomassie Blue. Lane *M* contains molecular-weight markers (Thermo Scientific Spectra Multicolor broad-range protein ladder; labelled in kDa). Lane 1 contains *VmoLac* protein.

condition consisting of 400 mM ammonium dihydrogen phosphate. Crystals grew in drops containing a 2:1 protein:reservoir ratio (Fig. 2).

4. Data collection

A cryoprotectant solution consisting of the crystallization solution supplemented with 30% (v/v) glycerol was added to the drop in order to exchange the solution containing the crystal. The crystal was subsequently mounted on a CryoLoop (Hampton Research) and flash-cooled in liquid nitrogen. X-ray diffraction intensities were collected on the ID29 beamline at the ESRF, Grenoble, France using a wavelength of 0.800 Å and a PILATUS 6M detector (DECTRIS, Switzerland) with 50 ms exposures. Diffraction data were collected using the fine-slicing method; individual frames consisted of 0.1° steps over a range of 100° (Fig. 3).

5. Results and conclusions

X-ray diffraction data were integrated and scaled using the *XDS* package (Kabsch, 1993; Table 1). The *VmoLac* crystals belonged to



Figure 2
Crystals of *VmoLac* (average dimensions of 100 × 50 × 40 μm).




Figure 3
A diffraction pattern from a crystal of *VmoLac*. The edge of the frame is at 2.0 Å resolution.

Table 1
Data-collection statistics.

Values in parentheses are for the last bin.

Beamline	ID29, ESRF
Wavelength (Å)	0.800
Detector	PILATUS 6M
Oscillation (°)	0.1
No. of frames	1000
Resolution (Å)	47.6–2.4 (2.5–2.4)
Space group	<i>P6₄</i>
Unit-cell parameters (Å)	<i>a</i> = <i>b</i> = 174.06, <i>c</i> = 61.32
No. of observed reflections	231397 (26406)
No. of unique reflections	41676 (4740)
Completeness (%)	99.6 (99.3)
<i>R</i> _{meas} † (%)	13.4 (63.5)
CC _{1/2} ‡	99.6 (81.8)
<i>I</i> /σ(<i>I</i>)	14.55 (3.22)
Multiplicity	5.55 (5.57)
Mosaicity (°)	0.201

† $R_{\text{meas}} = \sum_{hkl} \{N(hkl)/[N(hkl) - 1]\}^{1/2} \sum_i |I_i(hkl) - \langle I(hkl) \rangle| / \sum_{hkl} \sum_i I_i(hkl)$. ‡ CC_{1/2} is the intra-data-set correlation coefficient calculated from the percentage of correlation between intensities (*I*₁ and *I*₂) from random half data sets (Karplus & Diederichs, 2012): CC_{1/2} = Corr(*I*₁, *I*₂).

the hexagonal space group *P6₄*, with unit-cell parameters *a* = *b* = 174.06, *c* = 61.32 Å. *VmoLac* being a 35 kDa protein, the calculated Matthews coefficient (Matthews, 1968) suggests that between two and three monomers are present per asymmetric unit (3.78 and 2.52 Å³ Da⁻¹, corresponding to 67.49 and 51.24% solvent content, respectively). An initial molecular replacement was performed using *Phaser* (McCoy *et al.*, 2007) with the structure of *SsoPox* (52% sequence identity; PDB entry 2vc5; Elias *et al.*, 2008) from which the fragment 265–277 was deleted as a starting model. Two molecules were initially placed in the asymmetric unit (*R* = 47.25%, *R*_{free} = 48.1%). The initial solution was then submitted to *ARP/wARP* (Morris *et al.*, 2003) for automated model construction. After 50 cycles of *ARP/wARP* and 20 cycles of refinement using *REFMAC* (Murshudov *et al.*, 2011), the *R* and *R*_{free} factor values were 18.95 and 22.38%, respectively. The electron-density maps revealed that, with the exception of some rotamers and loop conformations, the model is near final. The asymmetric unit contains one homodimer of *VmoLac*. The construction, refinement and interpretation of the structure are in progress.

This research was supported by a grant to EC from Délégation Générale pour l'Armement (REI #2009 34 0045). GG and JH are supported by Délégation Générale pour l'Armement.

References

- Afriat, L., Roodveldt, C., Manco, G. & Tawfik, D. S. (2006). *Biochemistry*, **45**, 13677–13686.
- Afriat-Jurnou, L., Jackson, C. J. & Tawfik, D. S. (2012). *Biochemistry*, **51**, 6047–6055.
- Amara, N., Krom, B. P., Kaufmann, G. F. & Meijler, M. M. (2011). *Chem. Rev.* **111**, 195–208.
- Ben-David, M., Elias, M., Filippi, J. J., Duñach, E., Silman, I., Sussman, J. L. & Tawfik, D. S. (2012). *J. Mol. Biol.* **418**, 181–196.
- Benning, M. M., Kuo, J. M., Raushel, F. M. & Holden, H. M. (1994). *Biochemistry*, **33**, 15001–15007.
- Berg, S. van den, Löfdahl, P. A., Härd, T. & Berglund, H. (2006). *J. Biotechnol.* **121**, 291–298.
- Chow, J. Y., Xue, B., Lee, K. H., Tung, A., Wu, L., Robinson, R. C. & Yew, W. S. (2010). *J. Biol. Chem.* **285**, 40911–40920.
- Del Vecchio, P., Elias, M., Merone, L., Graziano, G., Dupuy, J., Mandrich, L., Carullo, P., Fournier, B., Rochu, D., Rossi, M., Masson, P., Chabriere, E. & Manco, G. (2009). *Extremophiles*, **13**, 461–470.
- Donarski, W. J., Dumas, D. P., Heitmeyer, D. P., Lewis, V. E. & Raushel, F. M. (1989). *Biochemistry*, **28**, 4650–4655.

- Dong, Y.-H., Wang, L.-H., Xu, J.-L., Zhang, H.-B., Zhang, X.-F. & Zhang, L.-H. (2001). *Nature (London)*, **411**, 813–817.
- Elias, M., Dupuy, J., Merone, L., Mandrich, L., Porzio, E., Moniot, S., Rochu, D., Lecomte, C., Rossi, M., Masson, P., Manco, G. & Chabriere, E. (2008). *J. Mol. Biol.* **379**, 1017–1028.
- Elias, M. & Tawfik, D. S. (2012). *J. Biol. Chem.* **287**, 11–20.
- Gotthard, G., Hiblot, J., Elias, M. & Chabrière, E. (2011). *Acta Cryst.* **F67**, 354–357.
- Gotthard, G., Hiblot, J., Gonzalez, D., Chabrière, E. & Elias, M. (2013). *Acta Cryst.* **F69**, 73–76.
- Gumerov, V. M., Mardanov, A. V., Beletsky, A. V., Prokofeva, M. I., Bonch-Osmolovskaya, E. A., Ravin, N. V. & Skryabin, K. G. (2011). *J. Bacteriol.* **193**, 2355–2356.
- Gupta, R. C. (2009). *Handbook of Toxicology of Chemical Warfare Agents*. London: Academic Press.
- Hawwa, R., Aikens, J., Turner, R. J., Santarsiero, B. D. & Mesecar, A. D. (2009). *Arch. Biochem. Biophys.* **488**, 109–120.
- Hawwa, R., Larsen, S. D., Ratia, K. & Mesecar, A. D. (2009). *J. Mol. Biol.* **393**, 36–57.
- Hiblot, J., Gotthard, G., Chabriere, E. & Elias, M. (2012a). *PLoS One*, **7**, e47028.
- Hiblot, J., Gotthard, G., Chabriere, E. & Elias, M. (2012b). *Sci. Rep.* **2**, 779.
- Jackson, C. J., Carr, P. D., Kim, H.-K., Liu, J.-W., Herrald, P., Mitić, N., Schenk, G., Smith, C. A. & Ollis, D. L. (2006). *Biochem. J.* **397**, 501–508.
- Jackson, C. J., Foo, J.-L., Tokuriki, N., Afriat, L., Carr, P. D., Kim, H.-K., Schenk, G., Tawfik, D. S. & Ollis, D. L. (2009). *Proc. Natl Acad. Sci. USA*, **106**, 21631–21636.
- Kabsch, W. (1993). *J. Appl. Cryst.* **26**, 795–800.
- Karplus, P. A. & Diederichs, K. (2012). *Science*, **336**, 1030–1033.
- LeJeune, K. E., Wild, J. R. & Russell, A. J. (1998). *Nature (London)*, **395**, 27–28.
- Masson, P., Carletti, E. & Nachon, F. (2009). *Protein Pept. Lett.* **16**, 1215–1224.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C. & Read, R. J. (2007). *J. Appl. Cryst.* **40**, 658–674.
- Merone, L., Mandrich, L., Porzio, E., Rossi, M., Müller, S., Reiter, G., Worek, F. & Manco, G. (2010). *Bioresour. Technol.* **101**, 9204–9212.
- Merone, L., Mandrich, L., Rossi, M. & Manco, G. (2005). *Extremophiles*, **9**, 297–305.
- Morris, R. J., Perrakis, A. & Lamzin, V. S. (2003). *Methods Enzymol.* **374**, 229–244.
- Murshudov, G. N., Skubák, P., Lebedev, A. A., Pannu, N. S., Steiner, R. A., Nicholls, R. A., Winn, M. D., Long, F. & Vagin, A. A. (2011). *Acta Cryst.* **D67**, 355–367.
- Omburo, G. A., Kuo, J. M., Mullins, L. S. & Raushel, F. M. (1992). *J. Biol. Chem.* **267**, 13278–13283.
- Seibert, C. M. & Raushel, F. M. (2005). *Biochemistry*, **44**, 6383–6391.
- Singh, B. K. (2009). *Nature Rev. Microbiol.* **7**, 156–164.
- Vyas, N. K., Nickitenko, A., Rastogi, V. K., Shah, S. S. & Quioco, F. A. (2010). *Biochemistry*, **49**, 547–559.
- Xue, B., Chow, J. Y., Baldansuren, A., Yap, L. L., Gan, Y. H., Dikanov, S. A., Robinson, R. C. & Yew, W. S. (2013). *Biochemistry*, **52**, 2359–2370.
- Zhang, Y., An, J., Ye, W., Yang, G., Qian, Z.-G., Chen, H.-F., Cui, L. & Feng, Y. (2012). *Appl. Environ. Microbiol.* **78**, 6647–6655.